

Kundurosaurus

Kundurosaurus is an extinct genus of saurolophine hadrosaurid dinosaur known from the Latest Cretaceous (probably Late Maastrichtian stage) of Amur Region, Far Eastern Russia. It contains a single species, ***Kundurosaurus nagorny***.^[1]

Contents
Description
Discovery
Classification
See also
References

Description



Dentaries AENM 2/846 (A-B) and AENM 2/902 (C-D).

Kundurosaurus is a saurolophine diagnosed by four autapomorphies, unique derived traits. It has a prominent and thick ridge on the lateral side of the nasal that borders caudally the circumnasal depression and invades the caudal plate of the nasal. Its caudal buttress of the proximal head of the scapula is oriented quite laterally, parallel to the pseudoacromial process. The preacetabular process of the ilium is straight and only moderately

deflected ventrally by an angle of 160°. It does not reach the level of the plane formed by the bases of the iliac and pubic peduncles. Finally, with *Kundurosaurus* the axis of the postacetabular process of the ilium is strongly twisted along its length, so that its lateral side progressively faces dorsolaterally.^[1]

Discovery

Kundurosaurus is known from holotype AENM 2/921, a partial, disarticulated skull, including a nearly complete braincase (AENM 2/921 1-2), two quadrates (3-4), squamosal (5), postorbital (6), frontal (7) and parietal (8) bones. The referred specimens are AENM 2/45-46, two jugals; AENM 2/83-84, 2/86, maxillae; AENM 2/57-58, nasals; AENM 2/48, postorbital; AENM 2/19, quadrate; AENM 2/121, 2/928 partial braincases; AENM 2/846, 2/902, dentaries; AENM 2/906, scapula; AENM 2/913, sternal; AENM 2/117, 2/903, 2/907-908, humeri; AENM 2/905, ulna; AENM 2/904, radius; AENM 2/922, nearly complete pelvic girdle and associated sacral elements. These were found at the same level as the holotype, but may belong to other individuals. All specimens are housed in the Amur Natural History Museum of the Institute of Geology and Nature Management, Russia.^[1]

Kundurosaurus was first described and named by Pascal Godefroit, Yuri L. Bolotsky and Pascaline Lauters in 2012 and the type species is *Kundurosaurus nagorny*. The generic name is derived from *Kundur*, the type and only known locality, and *sauros*, "lizard" in Ancient Greek. The specific name, *nagorny*, honors V.A. Nagorny from the Far Eastern Institute of

Kundurosaurus

Temporal range: Late Cretaceous, 67–66 Ma

PreЄ Є OS D C P T J K PgN

Restoration of the skull

Scientific classification

Kingdom: Animalia

Phylum: Chordata

Clade: Dinosauria

Order: †Ornithischia

Suborder: †Ornithopoda


Family: †Hadrosauridae

Subfamily: †Saurolophinae

Genus: †*Kundurosaurus*
Godefroit *et al.*, 2012

Type species

†***Kundurosaurus nagorny***
Godefroit *et al.*, 2012



Endocranial reconstruction of AENM 2/121 based on a CT scan.

All *Kundurosaurus* specimens were collected in the Kundur locality. The site belongs to the *Wodehouseia spinata* – *Aquilapollenites subtilis* palynozone, dating to the Maastrichtian stage, probably the Late Maastrichtian, of the Late Cretaceous period, about 67-66 million years ago. The Kundur site was discovered by Vladimir A. Nagorny in 1990. He collected fossil bones in a road section along the Chita – Khabarovsk highway near the village of Kundur and sent them to Yuri L. Bolotsky. Large-scale excavations started at Kundur in 1999. Besides the abundant *Olorotitan arharensis* material, it has yielded many disarticulated saurolophine specimens. All these specimens were assigned to *Kundurosaurus* because the describers considered the recovered material to be homogeneous, and suggested that there is no reason to believe that more than one single saurolophine taxon lived in the Kundur area by latest Cretaceous period.^[1]

Kundur is one of four rich dinosaur localities that have been discovered in the southeastern part ("Lower Zeya depression") of Zeya-Bureya sedimentary basin, eastern Asia: Jiayin and Wulaga localities are located in the Yuliangze Formation of northern Heilongjiang Province, China and Blagoveschensk and Kundur localities are located in the Udurchukan Formation of southern Amur Region, Russia. In each locality, the dinosaur fauna is largely dominated by lambeosaurine hadrosaurids (*Charonosaurus jiayinensis* and some non-diagnostic material of *Mandschurosaurus amurensis* from Jiayin, *Sahaliyana* from Wulaga, *Amurosaurus* from Blagoveschensk, and *Olorotitan* from Kundur),^[1] but the indeterminate hadrosaurid *Arkharavia*,^[2] from Kundur, and saurolophine (non-crested or solid-crested) hadrosaurids are also represented (*Saurolophus kryschtofovici* and other non-diagnostic material of *M. amurensis* from Jiayin, *Wulagasaurus* from Wulaga, *Kerberosaurus* from Blagoveschensk and *Kundurosaurus* from Kundur).^[1]


Classification

A phylogenetic analysis of saurolophines performed by Godefroit, Bolotsky & Lauters (2012) indicates that *Kundurosaurus* is nested within a clade including Edmontosaurini and Saurolophini, possibly as a sister-taxon of *Kerberosaurus*. It is based on the data matrix of Prieto-Márquez (2010), however Prieto-Márquez (2010) recovered Edmontosaurini as a sister-taxon of a monophyletic clade formed by Saurolophini and Kritosaurini while in Godefroit *et al.* (2012) the Edmontosaurini + Saurolophini clade is well supported and excludes Kritosaurini.^[1]

The position of *Kundurosaurus* within Edmontosaurini collapses when fragmentary taxa are excluded from the analysis. In the full analysis, *Kundurosaurus* is placed as the sister-taxon of *Kerberosaurus*, which is known from the same region. It may therefore be postulated that *K. nagorny* is a second species of the genus *Kerberosaurus*. This clade, however, is very weakly supported and synapomorphies uniting both taxa can only be found under optimization. Furthermore, although overlapping materials between the genera are limited to their partial skulls, according to Godefroit *et al.* (2012)



Location of hadrosaur localities in the Maastrichtian Zeya-Bureya Basin. Map of Heilongjiang, China, as well as the Amur Region of Russia. Blue dots represent the Yuliangze Formation, and red dots represent the Udurchukan Formation. Wulaga is home to *Wulagasaurus* and *Sahaliyana*, Jiayin to *Charonosaurus*, Kundur to *Kundurosaurus* and *Olorotitan*, and Blagoveschensk to *Kerberosaurus* and *Amurosaurus*.




Pelvic girdle of *Kundurosaurus*: left pubis AENM 2/922-5L (A), left ischium AENM 2/922-3L (B) and the autapomorphic left ilium AENM 2/922-7L (C) in lateral view.


Kundurosaurus can be differentiated from *Kerberosaurus* on the basis of the rostrocaudally longer and more robust dorsal maxillary process, more robust and more curved downwards nasal, much more robust and proportionally higher quadrate and the strong ridge extends obliquely along the lateral side of the exoccipital condyloid in *Kundurosaurus*. Additionally, the frontals of *Kerberosaurus* are particularly narrow and do not participate in the orbital margin, the rostral margin of the parietal is depressed around the contact area with the frontals, and *Kerberosaurus* has hook-like palatine process.^[1]

On the other hand, Xing *et al.* (2014) considered *Kundurosaurus nagorny* to be a junior synonym of *Kerberosaurus manakini* on the basis of their co-occurrence within the same formation and presence of shared characters in their skeletons.^[3]

The cladogram below follows Godefroit *et al.* (2012) analysis.^[1]




Scapula




Jugals



Right nasals





See also

- Timeline of hadrosaur research

References

1. Godefroit, P.; Bolotsky, Y. L.; Lauters, P. (2012). Joger, Ulrich (ed.). "A New Saurolophine Dinosaur from the Latest Cretaceous of Far Eastern Russia" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3364265>). *PLoS ONE*. **7** (5): e36849. doi:10.1371/journal.pone.0036849 (<https://doi.org/10.1371%2Fjournal.pone.0036849>). PMC 3364265 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3364265>). PMID 22666331 (<https://pubmed.ncbi.nlm.nih.gov/22666331>).
2. Godefroit, P. (2011). "Osteology and relationships of *Olorotitan arharensis*, a hollowcrested hadrosaurid dinosaur from the latest Cretaceous of Far Eastern Russia" (<https://doi.org/10.4202/app.2011.0051>). *Acta Palaeontologica Polonica*. doi:10.4202/app.2011.0051 (<https://doi.org/10.4202%2Fapp.2011.0051>).
3. Xing, Hai; Zhao, Xijin; Wang, Kebai; Li, Dunjing; Chen, Shuqing; Mallon, Jordan C; Zhang, Yanxia; Xu, Xing (2014). "Comparative osteology and phylogenetic relationship of *Edmontosaurus* and *Shantungosaurus* (Dinosauria: Hadrosauridae) from the Upper Cretaceous of North America and East Asia". *Acta Geologica Sinica-English Edition*. **88** (6): 1623–1652. doi:10.1111/1755-6724.12334 (<https://doi.org/10.1111%2F1755-6724.12334>).

Retrieved from "<https://en.wikipedia.org/w/index.php?title=Kundurosaurus&oldid=951380192>"

This page was last edited on 16 April 2020, at 21:45 (UTC).

Text is available under the [Creative Commons Attribution-ShareAlike License](#); additional terms may apply. By using this site, you agree to the [Terms of Use](#) and [Privacy Policy](#). Wikipedia® is a registered trademark of the [Wikimedia Foundation, Inc.](#), a non-profit organization.